

1.5T / 3T SCANNER EXAM PREFERENCES

(Updated 4/30/25)

1.5T	3T
• MRA Chest W/O (CIC, MID, QRY, CP MR1 only) • Pelvis Placenta Accreta • Spine Cervical • Spine Lumbar • Spine Thoracic 3T Contraindicated Patients, preferred at WLK or WMC. Also available at: • Fingers, GTN MR1 • Soft Tissue Neck, GTN MR1 or SW MR3 • Toes, GTN MR1	• Abdomen Liver lab for Hemochromatosis / Iron overload (WP) • Abdomen MRElastography (WP, MNR) • Brachial Plexus • Brain ARIA • Brain CSF Leak • Brain DBS Planning • Brain Perfusion • Brain Seizure (Adult/Pedi) • Brain Temporal Arteritis • Brain Vasculitis • Breasts • Enterography • Fingers • MRA Abdomen W/O (MPT only) • MRA Lower Extremity W/O (MPT only) • MSK Neurography • Multiple Sclerosis – UTHA Only (brain / orbits / cervical & thoracic spine) • Pelvis Anal Fistula • Pelvis Athletic Pubalgia • Pelvis Female for Cancer • Pelvis Neuro / Sacral Plexus • Pelvis Prostate • Pelvis Rectum • Pelvis Urethral Diverticulum • Skull Base / Face (lesion staging) • Soft Tissue Neck • Toes <i>Exams may be performed routinely at WLK or WMC.</i>