

Austin Radiological Association

MRI MSK Protocols

Adult 1.5T

Questions?

Last Update: 9/13/2024 8:15 AM

1.5T MSK Protocols

***1.5T Preferred Exams**

General Guidelines.....	4
1.5T and 3T Preferred Exams	6
Arthrogram.....	7
Ankle	8
Elbow	9
Hip	10
Knee.....	11
Shoulder	12
Wrist	13
Abdominal Wall	14
Chest Wall	15
Joint Contrast.....	16
Long Bone.....	17
Upper Extremity	18
Clavicle	18
Elbow	19
Hand	20
Pectoralis Tendon.....	21
Scapula	22
Shoulder	23
Sternum.....	24
Sternoclavicular Joints.....	25
Wrist	26
Lower Extremity.....	27

Achilles Tendon.....	28
Ankle / Hindfoot.....	29
Foot: Mid or Forefoot.....	30
Foot: Morton’s Neuroma or Contrast.....	31
Foot: Plantar Fibromatosis	32
Hip: Bilateral.....	33
Hip: Unilateral.....	35
Hip: Replacement	36
Knee.....	37
Knee: Metal.....	38
Knee: OtisMed Stryker.....	39
Knee: Visionaire, smith&nephew	40
Pelvis.....	41
Pelvis Contrast.....	42
Sacroiliac Joints.....	43
Tibia.....	44

General Guidelines

(Updated 9/13/24)

MSK

MSK	
General	<ul style="list-style-type: none">• NEVER hesitate to reach out to a radiologist for guidance!• All PEDI MSK arthritis exams should be performed using the standard adult protocols, unless specifically requested for pediatric radiologist to read• MSK Guidelines portal page reference<ul style="list-style-type: none">○ T1 & T2 weighted imaging is needed to accurately differentiate mass and infection (osteomyelitis, cellulitis, abscess)○ PDs are generally useful for tendons, ligaments, & joint spaces○ T2s are the most beneficial sequences and should be repeated for motion
Technique	<ul style="list-style-type: none">• Use “Weak” FS (Siemens) or “Classic” FS (GE) on all sequences with FS
Protocol	<ul style="list-style-type: none">• MSK <i>known</i> mass / infection (osteomyelitis, cellulitis, abscess) exams should be performed using T1 & T2 FS all 3 planes with appropriate T1 FS pre & post. Metastatic concerns do not require contrast<ul style="list-style-type: none">○ Always need<ul style="list-style-type: none">▪ <i>Known</i> mass / tumor exams - request prior x-ray study for comparison, if applicable▪ T1 FS pre – prior to contrast injection for direct comparison with post imaging▪ Generally short axis to the mass is the best option for pre/post imaging▪ T1 (generally Coronal) for ALL bone metastasis, infection, or fracture cases▪ If contrast is contraindicated a T1 FS pre does not need to be performed• Metal Protocols<ul style="list-style-type: none">○ Total joint replacements use T1 & IR series (TSE/FSE sequence types)○ Hardware such as pins, screws, etc. use standard protocols with metal reduction techniques○ Metal reduction techniques<ul style="list-style-type: none">▪ Bandwidth 400 Hz or more with signal compensation, fast RF mode▪ E-line (3T & Aera) WARP on, VAT 100%• Pelvis vs Hip

	<ul style="list-style-type: none"> ○ Hip – evaluation of cartilage & labrum ○ Pelvis – evaluation of fractures, soft tissue & cancer <ul style="list-style-type: none"> ▪ AVN, Osteonecrosis or history of “steroid use with hip pain” ▪ Cancer (mass, tumor), metastasis, myeloma ● Inflammatory Arthritis (rheumatoid arthritis, psoriatic arthritis, juvenile rheumatoid arthritis, reactive arthritis, gout, CPPD [calcium pyrophosphate deposition disease], or septic arthritis) <ul style="list-style-type: none"> ○ The entire hand and entire foot must be scanned for arthritis cases without any other indications such as a tear or mass <ul style="list-style-type: none"> ▪ Contrast is needed to differentiate joint fluid from synovitis ○ If a hand and wrist exam are ordered with a diagnosis of arthritis without any other indications, the wrist and hand need to be scanned together in the same FOV ● History of pigmented villonodular synovitis (PVNS) <ul style="list-style-type: none"> ○ Add T1 Coronal for all joint studies
<p style="text-align: center;">Contrast</p>	<ul style="list-style-type: none"> ● Delayed contrast required for all exams, perform at least 1 T2 weighted image immediately after injection to allow for sufficient uptake time. ● Two planes post contrast are always needed on MSK exams ● Not required for marrow disease or lesions (mass/tumor) unless in rare cases it extends into the soft tissues ● X-ray / CT abdomen and pelvis imaging must be performed prior to MR contrast exams. ● DatScans must be performed prior to MR contrast exams.

1.5T and 3T Preferred Exams

1.5T	3T
•	<ul style="list-style-type: none">• Fingers• Pelvis Athletic Pubalgia• Toes
Hand and foot exam codes are closed on 1.5T scanners to avoid incorrect scheduling of finger and toe exams. See 3T MSK Protocols	

Arthrogram




GENERAL GUIDELINES

Failed Exam	<ul style="list-style-type: none">• Perform routine joint protocol for failed exams, i.e., no contrast within the joint space. Radiologist will interpret study or determine if patient needs to return.
Indirect Arthrogram of Any Joint	<ul style="list-style-type: none">• Requires Radiologist consultation / approval• Inject IV gadolinium• Instruct patient to move the affected joint for 15 minutes in rotation, abduction, & adduction prior to scanning• Follow the applicable routine Arthrogram protocol
Available Locations	<ul style="list-style-type: none">• QRY• SW (MR1, MR2, MR3)

Ankle

Arthrogram protocols must include tech notes regarding the patient's pain post arthrogram "fluoro" procedure.




- Is the pain the same, improved or worsened since the procedure?

SEQUENCE	FOV (mm)	SLICE (mm)	COMMENTS	IMAGES
T1 Ax T1 FS Ax	~140	3.5 x 1 ~24 slices		
T1 FS Cor	~140	3.5 x 1 ~32 slices	Perpendicular to lateral/medial malleoli	
T1 FS Sag T2 FS Sag	~140	3 x 1 ~21 slices	Parallel to lateral/medial malleoli	

Elbow

Arthrogram protocols must include tech notes regarding the patient's pain post arthrogram "fluoro" procedure.

- Is the pain the same, improved or worsened since the procedure?

SEQUENCE	FOV (mm)	SLICE (mm)	COMMENTS	IMAGES
T1 Ax T1 FS Ax	120	4 x 1 ~24 slices	Include the radial tuberosity to include the bicep tendon attachment	
T1 FS Cor T2 FS Cor	~140	3 x 0 ~30 slices	<ul style="list-style-type: none"> • Parallel to the humeral epicondyles • Do not use a localizer to set the angle of the coronal series 	
T1 FS Sag	~140	3 x 1 ~21 slices	<ul style="list-style-type: none"> • Perpendicular to the humeral epicondyles • Do not use a localizer to position the sagittal series 	




Hip

(Updated 5/21/21)

Consider MSK Pelvis protocol with history of arthroplasty / total joint replacement, consult with MSK Radiologist.

Position:

- Tape or strap feet with internal rotation, note patient's inability to do so
- Legs need to be as flat as possible

SEQUENCE	FOV (mm)	SLICE (mm)	COMMENTS	IMAGES
T1 Cor Bilat STIR Cor Bilat	~360	4 x 0 28 slices	Parallel to the anterior surfaces of the femoral heads	
T1 Ax Bilat (TE 12 – 20) T2 FS Ax Bilat (STIR Espree or suboptimal FS)	~360	5 x 1 ~36 slices	<ul style="list-style-type: none"> • Parallel to the superior surfaces of the femoral heads • Include from above sacrum through the hamstring attachment on the ischium 	
PD FS Ob l Ax (TE 40 – 45) PD FS Cor PD FS Sag	~200	4 x 1 ~24 slices	Ax: above acetabulum through ischium	
Contrast, if needed T1 FS Cor Pre / Post	~360	4 x 0 28 slices	Perform T2 weighted image immediately after injection to allow for contrast uptake.	Copies to T1 Cor
T1 FS Ax Post	~360	5 x 1 ~36 slices		Copies to T1 Ax

- Include the hamstring attachment on the ischial tuberosity on all sequences

Tech Notes:




Arthrogram protocols must include tech notes regarding the patient's pain post arthrogram "fluoro" procedure.

- Is the pain the same, improved or worsened since the procedure?






Knee

Arthrogram protocols must include tech notes regarding the patient's pain post arthrogram "fluoro" procedure.

- Is the pain the same, improved or worsened since the procedure?

SEQUENCE	FOV (mm)	SLICE (mm)	COMMENTS	IMAGES
T1 FS Ax	~140	4 x 1 ~24 slices		
T1 FS Sag PD Sag	~140	3 x 1 ~24 slices	Parallel to the outer surface of the lateral femoral condyle	
T1 FS Cor T2 FS Cor	~140	3 x 1 ~24 slices	Parallel to the posterior surfaces of the femoral condyles	




Shoulder

Position				
<ul style="list-style-type: none"> • Supine with affected arm at side in external rotation (palm directed upward) • Place sandbag on palm if necessary to help the patient maintain this position 				
SEQUENCE	FOV (mm)	SLICE (mm)	COMMENTS	IMAGES
T1 FS Ax T2 FS Ax	~140	4 x 1 ~22 slices	Include AC joint	
T1 FS Cor T2 FS Cor	~140	3 x 1 ~18 slices	Perpendicular to the glenoid fossa	
T1 Sag	~140	4 x 1	Parallel to the glenoid fossa	
T2 FS Sag		3 x 1 ~24 slices		
Localizer			<ul style="list-style-type: none"> • Position the affected arm above the patient's head with the elbow flexed and the palm of the hand under the patient's head. • Send the localizer to PACS 	
T1 FS ABER	~140	3 x 0.75 ~24 slices	<ul style="list-style-type: none"> • Parallel to the long axis of the humerus and centered to the joint 	
Arthrogram protocols must include tech notes regarding the patient's pain post arthrogram "fluoro" procedure. <ul style="list-style-type: none"> • Is the pain the same, improved or worsened since the procedure? 				

Wrist

Arthrogram protocols must include tech notes regarding the patient's pain post arthrogram "fluoro" procedure.




- Is the pain the same, improved or worsened since the procedure?

SEQUENCE	FOV (mm)	SLICE (mm)	COMMENTS	IMAGES
T2 FS Ax	~100	3 x 0 ~24 slices		
T1 Cor T1 FS Cor T2 FS Cor	~100	3 x 0 ~14 slices		
T1 FS Sag	~100	3 x 0 ~24 slices		

Abdominal Wall

(mass, focal pain, fracture of rib, costochondral, costochondritis, specific area of interest)




(Updated 10/15/20)

<ul style="list-style-type: none"> • Tailor coverage to area of interest • Run phase direction & position patient supine/prone to best minimize motion artifact. • Perform as breath-hold as needed to minimize motion artifacts. • Do no perform at ARA RCP MR1 GE, SM, WLK, WMC 				
SEQUENCE	FOV (mm)	SLICE (mm)	COMMENTS	IMAGES
T1 Ax STIR Ax	240	5 x 1.3		
T1 Cor STIR Cor	240	5 x 1.3		
T1 Sag STIR Sag	240	4 x 0		
<i>Contrast, if needed</i> T1 FS AX Pre / Post Additional plane post			Perform T2 weighted image immediately after injection to allow for contrast uptake.	
Exam Code: <ul style="list-style-type: none"> • MRABWCSMSK – MRI Abdominal Wall With and Without Contrast – MSK • MRABWSMSK – MRI Abdominal Wall Without Contrast - MSK 				

Chest Wall

(mass, focal pain, fracture of rib, costochondral, costochondritis, specific area of interest)

(Updated 10/15/20)

<ul style="list-style-type: none"> • If FS is bad, run STIR • Tailor coverage to area of interest • Run phase direction & position patient supine/prone to best minimize motion artifact • Perform as breath-hold as needed to minimize motion artifacts • If contrast pick a second plane post that shows lesion the best 				
SEQUENCE	FOV (mm)	SLICE (mm)	COMMENTS	IMAGES
T1 Ax STIR Ax	250	4 x 0.5		
T1 Cor STIR Cor	250	5 x 1		
T1 Sag STIR Sag	250	5 x 1		
<i>Contrast, if needed</i> T1 FS AX Pre / Post Additional plane post			Perform T2 weighted image immediately after injection to allow for contrast uptake.	

Joint Contrast

(synovitis, rheumatoid arthritis, inflammatory arthritis, or inflammatory arthropathy)

- Tailor FOV to body part
- Perform T2 weighted image immediately after injection to allow for contrast uptake.




ADDITIONAL SEQUENCES	COMMENTS
T1 FS Ax Pre / Post T1 FS Sag Post	Ankle, Foot
<u>Follow Hip: Unilateral protocol</u>	Hip
T1 FS Cor Pre / Post T1 FS Ax Post	Shoulder, Hand, Wrist
T1 FS Cor Pre / Post T1 FS Sag Post	Elbow, Knee

Long Bone

- Place skin marker to indicate area of concern
- Partial FOVs – focal pain, mass; must include either the proximal or distal joint
- Full FOVs – general pain, myositis, cellulitis; upper/lower as needed to include full long bone

Position

- Humerus: anatomical with palm face up
- Forearm: arm down by side or superman with palm face up
- Femur: anatomical with toes pointing up
- Tib/Fib: anatomical with toes pointing up

SEQUENCE	FOV (mm)	SLICE (mm)	COMMENTS	IMAGES
T1 Ax STIR Ax	140 – 200	5 x 1	Humerus: Forearm: parallel to radius / ulna Femur: Tib/Fib:	
T1 Cor STIR Cor	200 - 300	6 x 1	Humerus: Forearm: parallel to radius / ulna Femur: parallel to femur Tib/Fib:	
T1 Sag STIR Sag	200 – 300	6 x 1		
Contrast: mass, infection, or post-op, may consult with radiologist for appropriate planes <i>Contrast not required for metastatic disease</i>				
T1 FS Ax Pre T1 FS Ax Post Additional plane post: T1 FS Cor Post T1 FS Sag Post			Perform T2 weighted image immediately after injection to allow for contrast uptake.	

Upper Extremity

Clavicle

- Position patient Prone
- All planes are orthogonal to body, do not angle with clavicle
- Include opposite SC joint in all series
- Perform STIR for suboptimal FS






SEQUENCE	FOV (mm)	SLICE (mm)	COMMENTS	IMAGES
T1 Ax T2 FS Ax	~200	3 x 0		
T1 Cor T2 FS Cor	~200	3 x 0		
T1 Sag T2 FS Sag	~200	4x1		

Elbow

(biceps tendon tear)

(Updated 6/15/22)




- See shoulder positioning aid for external rotation of humeral head

SEQUENCE	FOV (mm)	SLICE (mm)	COMMENTS	IMAGES
PD FS Ax T2 FS Ax	~140	4 x 1 ~24 slices	<ul style="list-style-type: none"> • Include radial tuberosity for bicep tendon attachment • Adjust in plane rotation for correct anatomical positioning 	
PD Cor T2 FS Cor	~140	3 x 0 ~30 slices	Position off PD FS Ax, parallel to humeral epicondyles	
T1 Sag T2 FS Sag / STIR	~140	3 x 1 ~21 slices	<ul style="list-style-type: none"> • Position off PD FS Ax, perpendicular to humeral epicondyles 	
T2 FS Long Axis <i>FABS View, optional Dr. Daniel Stewart, TX Orthopedics</i>	~130	3 x 0.5 ~24 slices	<ul style="list-style-type: none"> • Position elbow within shoulder coil in 90° flexion with forearm and palm in lateral position. • <u>FABS View Positioning</u> – Shoulder Coil 	 



Hand (pain)

- Tailor coverage to area of interest, 3T often performs as partial hand
- Always place marker to indicate area of interest


SEQUENCE	FOV (mm)	SLICE (mm)	COMMENTS	IMAGES
T1 Ax T2 FS Ax	120	4 x 1 ~22 slices	<ul style="list-style-type: none"> • Oblique to obtain true axial 	
T1 Cor T2 FS Cor	220	2 x 0.5 ~23 slices	<ul style="list-style-type: none"> • Oblique to obtain true coronal 	
PD Sag T2 FS Sag	220	4 x 1 ~24 slices	<ul style="list-style-type: none"> • Oblique to obtain true sagittal • Include adjacent 2nd – 5th finger for comparison, do not use thumb 	

Pectoralis Tendon

(Updated 6/5/23)

Position:

- Patient prone with the affected arm as close to the patient's body as possible to achieve the 20cm FOV
- Affected side as close to isocenter as possible to ensure quality imaging through the humerus laterally






SEQUENCE	FOV (mm)	SLICE (mm)	COMMENTS	IMAGES
PD Ax STIR Ax	~200	4 x 0.5 ~40 slices	Scan affected side only	
T1 Cor T2 FS Cor	~200	4 x 0 ~24 slices	Parallel to the pectoral muscle/tendon and through the humeral shaft	
T1 Sag T2 FS Sag	~200	4 x 0.5 ~30 slices	Perpendicular to pectoral muscle/tendon and through humeral shaft	
STIR Ax Bilat	~400	5 x 1 ~32 slices	<ul style="list-style-type: none"> • Including both sides with arms at the patient's side. • Espree magnets, right / left sides should be scanned separately due to the bore FOV limitations. 	

- The small FOVs are designed to show tears of the pectoral tendon at its insertion on the humerus or at the junction of the pectoral muscle and pectoral tendon. A 20cm FOV should be achievable on most patients. Do not need to include the entire pectoral muscle.
 - L>R: mid pectoral muscle through humerus
 - I>S: top of the pectoral muscle through the deltoid tuberosity of the humerus
 - P>A: posterior to the humerus through the pectoral muscle anteriorly
- The large FOV STIR Axial sequence will show muscle tears that aren't seen on the small FOV sequences
 - Position patient prone with both arms down by side

Scapula






(Updated 2/8/07)

SEQUENCE	FOV (mm)	SLICE (mm)	COMMENTS	IMAGES
T1 Ax T2 FS Ax	220	6 x 1 ~24 slices	Oblique to obtain true axial	
T1 Cor STIR Cor	240	4 x 1 ~20 slices	Oblique to obtain true coronal	
T1 Sag T2 FS Sag	240	6 x 1 ~25 slices	Oblique to obtain true sagittal	

Shoulder

Position:

- Affected arm at side with external rotation, palm facing upward
- Place sandbag on palm if necessary to help the patient maintain this position




SEQUENCE	FOV (mm)	SLICE (mm)	COMMENTS	IMAGES
PD FS Ax (TE 40 – 45) STIR Ax	150	3 x 0 ~22 slices	<ul style="list-style-type: none"> • Scan through the AC joint • Sag loc – slices parallel with the acromion & coracoid • Cor loc – slices perpendicular to the glenoid 	
PD Cor T2 FS Cor	140	3 x 1 ~18 slices	<ul style="list-style-type: none"> • Slices need to be perpendicular to the glenoid fossa • Sag loc – position slices perpendicular to scapular spine & coracoid • PD FS Ax – position slices parallel with supraspinatus tendon 	
T1 Sag	140	4 x 1	<p>T1 Sag: Include 6cm medial to the glenoid fossa</p> <ul style="list-style-type: none"> • Slices need to be parallel to the glenoid fossa 	
T2 FS Sag	140	3 x 1	<ul style="list-style-type: none"> • PD FS Ax – position slices perpendicular to the supraspinatus tendon • PD Cor – position slices parallel with the glenoid fossa • Verify in plane rotation with sag localizer is orthogonal 	
T1 Ax T2 FS Ax <i>Dr. Douglas, Austin Sports Medicine</i>	150	5 x 1	<ul style="list-style-type: none"> • Remove shoulder coil and use torso • Include shoulder joint through mid humerus 	

Sternum

Position

- Prone when possible
- Use of breast coil is encouraged

- Perform STIR for suboptimal FS
- Orient phase direction that best minimizes artifact




SEQUENCE	FOV (mm)	SLICE (mm)	COMMENTS	IMAGES
T1 Ax T2 FS Ax	180	3 x 0		
T1 Cor T2 FS Cor	180	3 x 0 ~18 slices		
T1 Sag T2 FS Sag	180	3 x 0 ~18 slices		

Sternoclavicular Joints

(Updated 5/29/08)




Position

- Prone when possible
- Perform STIR for suboptimal FS
- Orient phase direction that best minimizes artifact

SEQUENCE	FOV (mm)	SLICE (mm)	COMMENTS	IMAGES
T1 Ax T2 FS Ax	~180	3 x 0 ~32 slices		
T1 Cor T2 FS Cor	~180	3 x 0 ~18 slices		
T1 Sag T2 FS Sag	~180	3 x 0 ~18 slices		

Wrist





- Perform T2 weighted image immediately after injection to allow for contrast uptake.

SEQUENCE	FOV (mm)	SLICE (mm)	COMMENTS	IMAGES
PD Ax PD FS Ax (TE 40 – 45)	100 256 x 167	3 x 0 ~24 slices		
T2 FS Cor PD Cor	100 256 x 167	3 x 0 ~21 slices		
T2 3D Cor	120	1.5 x 0 ~36 slices		Copies center to T2 FS Cor
T1 Cor, <i>optional</i>	100	3 x 0 ~21 slices	Add for history of trauma or bright bone on T2	Copies to T2 FS Cor
STIR Sag	100 256 x 167	3 x 0 ~24 slices		
Contrast: pain				
T1 FS Cor Pre T1 FS Cor Post T1 FS Ax Post				
Contrast: mass protocol (T1, T2 FS protocol)				
T1 FS Ax Pre T1 FS Ax Post Additional plane to best demonstrates mass				
Contrast: inflammatory process				
Follow <u>JOINT CONTRAST</u> protocol				
Not performed on WLK MR1 Symphony				

Lower Extremity

Achilles Tendon

(Updated 12/13/18)





• Include all tarsal bones on all sequences				
SEQUENCE	FOV (mm)	SLICE (mm)	COMMENTS	IMAGES
PD Ax T2 FS Ax	~150	3.5 x 1 ~36 slices		
T1 Cor T2 FS Cor	~160	3.5 x 1 ~32 slices	Perpendicular to lateral/medial malleoli, see talar dome	
T1 Sag STIR Sag	~160	3 x 1 ~21 slices	Parallel to lateral/medial malleoli, see talar dome	
T2 FS Sag	250	3 x 1 ~21 slices	<ul style="list-style-type: none"> • Parallel with Achilles tendon • Include mid-calf through calcaneus 	

Increase FOV on sagittal series if the Achilles is torn and retracted outside of 160 FOV.


Ankle / Hindfoot

(pain, plantar fasciitis, posterior tibial tendon pathology)

- Include all tarsal bones on all sequences

SEQUENCE	FOV (mm)	SLICE (mm)	COMMENTS	IMAGES
PD Ax T2 FS Ax	~140	3.5 x 1 ~24 slices		
PD Obl Ax	~140	3.5 x 1 ~32 slices	Use angle of 35-45 degrees	
T1 Cor T2 FS Cor	~140	3.5 x 1 ~32 slices	Perpendicular to lateral/medial malleoli, see talar dome	
T1 Sag STIR Sag	~140	3 x 1 ~21 slices	Parallel to lateral/medial malleoli, see talar dome	

- Must include Lisfranc joint in all planes
- If the referral states plantar fascia fibromatosis, verify the diagnosis since it is usually plantar fasciitis not fibromatosis
- Fibromatosis requires a 200 FOV in all 3 planes to include all the foot except the toes



Foot: Mid or Forefoot (metatarsal pain, swelling or mass)


• Include all tarsal bones on all sequences				
SEQUENCE	FOV (mm)	SLICE (mm)	COMMENTS	IMAGES
T1 Ax T2 FS Ax	140	3.5 x 0.5 ~24 slice		
T1 Cor T2 FS Cor	140	4 x 1 ~34 - 42 slices	Perpendicular to lateral/medial malleoli, see talar dome	
T1 Sag STIR Sag	140	3 x 1 ~23 slices	Angle to specific metatarsal as needed	
Must include Lisfranc joint in all planes				
<p>Forefoot: Morton's Neuroma, includes indications of interspace evaluation</p> <p>Toe: osteomyelitis</p> <p>Whole foot: plantar nerve lesion</p>				

Foot: Morton's Neuroma or Contrast

(interspace, metatarsal pain, swelling or mass)

Updated 11/30/21




- Morton's Neuroma – FOV to include mid-foot through end of toes, include all tarsal bones on all sequences
- Contrast – consult radiologist for coverage as needed



SEQUENCE	FOV (mm)	SLICE (mm)	COMMENTS	IMAGES
T1 Ax T2 FS Dixon Ax	140	3 x 1 ~24 slice		
T1 Cor T2 FS Dixon Cor	140	3 x 1 ~34 slices	Perpendicular to lateral/medial malleoli, see talar dome	
T1 Sag STIR Sag	140	2 x 0.5 ~33 slices	Angle to specific metatarsal as needed	
T1 FS Dixon Cor Pre <i>Administer contrast</i>			Perform T2 weighted image immediately after injection to allow for contrast uptake.	Copies to T1 Cor
T1 FS Dixon Cor Post				
T1 FS Dixon Ax Post				Copies to T1 Ax

Foot: Plantar Fibromatosis




- Include all the foot except the toes

SEQUENCE	FOV (mm)	SLICE (mm)	COMMENTS	IMAGES
T1 Ax T2 FS Dixon Ax	200	3 x 1		
T1 Cor T2 FS Dixon Cor	140	3 x 1 ~50 slices	Perpendicular to lateral/medial malleoli, see talar dome	
T1 Sag T2 FS Dixon Sag	200	2 x 0.5 ~33 slices	Parallel to lateral/medial malleoli, see talar dome	



- If the referral states plantar fascia fibromatosis, verify the diagnosis since it is usually plantar fasciitis not fibromatosis
- Fibromatosis is a rare, benign, hyperproliferative fibrous tissue disorder resulting in the formation of nodules along the plantar fascia. It is the thickening of the plantar fascia

Hip: Bilateral

(Updated 11/6/20)

Perform for partial and total hip arthroplasty				
Position:				
<ul style="list-style-type: none"> • Tape or strap feet with internal rotation • Legs need to be as flat as possible 				
SEQUENCE	FOV (mm)	SLICE (mm)	COMMENTS	IMAGES
T1 Cor Bilat STIR Cor Bilat	~360	4 x 0 28 slices	Parallel to the anterior surfaces of the femoral heads	
T1 Ax Bilat (TE 12 – 20) T2 FS Ax Bilat (STIR Espree or suboptimal FS)	~360	5 x 1 ~36 slices	<ul style="list-style-type: none"> • Parallel to the superior surfaces of the femoral heads • Include from above sacrum through the hamstring attachment on the ischium 	
RT PD FS Obl Ax (TE 40 – 45) LT PD FS Obl Ax (TE 40 – 45)	~200	4 X 1 ~24 slices	Ax: above acetabulum through ischium	

- continued to next page -




RT PD FS Cor LT PD FS Cor	~200	4 X 1 ~20 slices		
RT PD FS Sag LT PD FS Sag	~200	4 X 1 ~26 slices		
Contrast: osteomyelitis, mass <i>Contrast not required for metastatic disease</i>				
T1 FS Cor Pre / Post	~360	4 x 0 28 slices	Perform T2 weighted image immediately after injection to allow for contrast uptake.	Copies to T1 Cor
T1 FS Ax Post	~360	5 x 1 ~36 slices		Copies to T1 Ax
<ul style="list-style-type: none"> • Include the hamstring attachment on the ischial tuberosity on all sequences <p>Tech Notes: Document patient's inability to internally rotate</p> <p>Send to PACS: Bilateral sequences to both folders</p>				

Hip: Unilateral

Consider MSK Pelvis protocol with history of arthroplasty / total joint replacement, consult with MSK Radiologist.

Position:

- Tape or strap feet with internal rotation
- Legs need to be as flat as possible

SEQUENCE	FOV (mm)	SLICE (mm)	COMMENTS	IMAGES
T1 Cor Bilat STIR Cor Bilat	~360	4 x 0 28 slices	Parallel to the anterior surfaces of the femoral heads	
T1 Ax Bilat (TE 12 – 20) T2 FS Ax Bilat (STIR Espree or suboptimal FS)	~360	5 x 1 ~36 slices	<ul style="list-style-type: none"> • Parallel to the superior surfaces of the femoral heads • Include from above sacrum through the hamstring attachment on the ischium 	
PD FS Ob l Ax (TE 40 – 45) PD FS Cor PD FS Sag	~200	4 X 1 ~24 slices	Ax: above acetabulum through ischium	
Contrast, if needed T1 FS Cor Pre / Post	~360	4 x 0 28 slices	<ul style="list-style-type: none"> • Consult radiologist for possible MSK pelvis if ordered for METS, lesion, mass • Perform T2 weighted image immediately after injection to allow for contrast uptake. 	Copies to T1 Cor
T1 FS Ax Post	~360	5 x 1 ~36 slices		Copies to T1 Ax

- Include the hamstring attachment on the ischial tuberosity on all sequences

Tech Notes:

Document patient's inability to internally rotate




Hip: Replacement

(DEPUY hip prosthesis recall)

Perform for total hip arthroplasty.

Position:

- Tape or strap feet with internal rotation
- Legs need to be as flat as possible

SEQUENCE	FOV (mm)	SLICE (mm)	COMMENTS	IMAGES
T1 Cor Bilat STIR Cor Bilat	~360	6 x 1	Cover entire boney pelvis	
T1 Ax Bilat STIR Ax	~360	5 x 1 ~36 slices	Cover entire boney pelvis	
T1 Sag	~360	6 x 1	Cover entire boney pelvis Espree – Cover as much as possible without orange lines	




Use Fast Spin Echo (FSE) sequences & a high bandwidth in order to reduce the artifact from the hip prosthesis. Siemens – up to 400 hz/pixel, GE – up to 64khz.

Knee

(Updated 10/3/23)

For indications of quad tendon:




- Torso coil
- FOV 17 CM for Sag / Cor – include patellar tendon attachment & above

SEQUENCE	FOV (mm)	SLICE (mm)	COMMENTS	IMAGES
PD FS Ax (TE 40 – 45)	140	4 x 1 ~24 slices		
PD Sag (TE 15, not above 20) PD FS Sag (TE 40 – 45)	140	3 x 1 ~24 slices	Parallel to the outer surface of the lateral condyle	
PD Cor (TE 15, not above 20) T2 FS Cor	140	3 x 1 ~24 slices	Parallel to the posterior surfaces of the femoral condyles	
T1 Cor, <i>optional</i>			<ul style="list-style-type: none"> • Add for history of trauma or bright bone on T2 • Add for PVNS (pigmented villonodular synovitis) 	Copies to PD Cor

Knee: Metal

(pin / screw vs total replacement)

(updated 4/13/2023)

Select protocol based on hardware, pin/screw vs total joint					
PIN / SCREWS		TOTAL JOINT		COMMENTS	IMAGES
SEQUENCE	PARAMETERS	SEQUENCE	PARAMETERS		
PD FS IR Ax	140 FOV 4 x 1 mm ~24 slices	T1 Ax IR Ax	150 FOV 4 x 1.3 mm		
PS Sag PD FS IR Sag	140 FOV 3 x 1 mm ~24 slices	T1 Sag IR Sag	150 FOV 4 x 1.3 mm	Parallel to the outer surface of the lateral condyle	
PD Cor T2 FS Cor	140 FOV 3 x 1 mm ~24 slices	T1 Cor IR Cor	150 FOV 4 x 1.3 mm	Parallel to the posterior surfaces of the femoral condyles	
T1 Cor, <i>optional</i>				<ul style="list-style-type: none"> • Add for history of trauma or bright bone on T2 • Add for PVNS (pigmented villonodular synovitis) 	Copies to PD Cor

Knee: OtisMed Stryker

Only perform at MPT MR2, 1.5T Espree	
SEQUENCE	REFERENCE
PD Sag	<u>Quick Reference</u> <u>Graphic Instructions</u> <u>Imaging Technique Training</u>
Body Coil	<u>Instructions</u>

Knee: Visionaire, smith&nephew

Available locations:

- 1.5T: MPT MR2 Espree
- 3T: Kyle, WP

Exam Codes: MRKNSVR, MRKNSVL

SEQUENCE	REFERENCE
Protocol	<u>Siemens Espree 1.5T</u>
3D Distortion Correction	<u>Instructions</u>

Pelvis

(inguinal hernia, coccyx or sacrococcygeal, hip for AVN)

See Pedi protocol for 0-17 y/o or small adults




Consider hip protocol if the diagnosis or patient history (particularly when the patient is under the age of 60) indicates possible hip pathology such as labral tear, cartilage abnormality, consult with MSK Radiologist.

Prep:

- Empty bladder

Position:

- Tape or strap feet with internal rotation
- Legs need to be as flat as possible

SEQUENCE	FOV (mm)	SLICE (mm)	COMMENTS	IMAGES
T1 Ax T2 FS Ax (STIR for Espree)	~360 (minimum FOV to include entire boney pelvis)	6 x 1 ~36 slices	Parallel to the superior surfaces of the femoral heads	
T1 Cor STIR Cor	~360 (minimum FOV to include entire boney pelvis)	5 x 1 ~32 slices		
T2 FS Sag T1 Sag	~300	6 x 1 (include entire boney pelvis)	T2 FS Sag: include entire boney pelvis T1 Sag: only include for sacrum/coccyx fracture	

- Include the hamstring attachment on the ischial tuberosity on all sequences

Tech Notes:

Document patient's inability to internally rotate

Pelvis Contrast

(sacroiliitis, arthritis, mass)




See Pedi protocol for 0-17 y/o or small adults

Prep:

- Empty bladder

Position:

- Tape or strap feet with internal rotation
- Legs need to be as flat as possible

SEQUENCE	FOV (mm)	SLICE (mm)	COMMENTS	IMAGES
T1 Ax T2 FS Ax (STIR for Espree)	~360 (minimum FOV to include entire boney pelvis)	6 x 1 ~36 slices	Parallel to the superior surfaces of the femoral heads	
T1 Cor STIR Cor	~360 (minimum FOV to include entire boney pelvis)	5 x 1 ~32 slices		
T2 FS Sag	~300	6 x 1 (include entire boney pelvis)	T2 FS Sag: include entire boney pelvis T1 Sag: only include for sacrum/coccyx fracture	
T1 Sag				
Contrast: sacroiliitis, arthritis				
T1 FS Cor Pre <i>Administer contrast</i> T1 FS Cor Post T1 FS Ax Post			Perform T2 weighted image immediately after injection to allow for contrast uptake.	
Contrast: mass <i>Contrast not required for metastatic disease</i>				
T1 FS Ax Pre <i>Administer contrast</i> T1 FS Ax Post T1 FS Cor Post			Perform T2 weighted image immediately after injection to allow for contrast uptake.	

Sacroiliac Joints

(Updated 1/6/23)

Only perform if exam is specifically ordered as SI Joint protocol.





Prep:

- Empty bladder

Position:

- Tape or strap feet with internal rotation
- Legs need to be as flat as possible

- If not administering contrast, do not perform T1 FS Cor pre

SEQUENCE	FOV (mm)	SLICE (mm)	COMMENTS	IMAGES
T2 FS Ax	~360 (minimum FOV to include entire boney pelvis)	6 x 1	Parallel to the superior surfaces of the femoral heads	
T1 Sag	~240	5 x 1	Include sacrum only	
T1 Ax Hires T2 FS Ax Hires	~240	5 x 1 ~30 slices	Perpendicular to Sacrum	
T1 Cor STIR Cor <i>Administer contrast, if needed</i> T1 FS Cor Pre / Post T1 FS Ax post (copies T1 Ax hires)	~240	5 x 1	<ul style="list-style-type: none"> • Parallel with Sacrum • Extend coverage to include cyst or posterior soft tissue, as needed • Perform T2 weighted image immediately after injection to allow for contrast uptake. 	
Send to PACS: Localizers				




Tibia

(Updated 5/1/24)

- Place skin marker to indicate area of concern
- Partial FOVs – focal pain, mass; must include either the proximal or distal joint
- Full FOVs – general pain, myositis, cellulitis; upper/lower as needed to include full long bone

Position

- Tib/Fib: anatomical with toes pointing up

SEQUENCE	FOV (mm)	SLICE (mm)	COMMENTS	IMAGES
T1 Ax STIR Ax	140 – 200	5 x 1		
T1 Cor STIR Cor	200 - 300	6 x 1	Tib/Fib: perpendicular to the tibiotalar joint	
T1 Sag STIR Sag	200 – 300	6 x 1	Tib/Fib: parallel to the tibiotalar joint	
Contrast: mass, infection, or post-op, may consult with radiologist for appropriate planes <i>Contrast not required for metastatic disease</i>				
T1 FS Ax Pre T1 FS Ax Post Additional plane post: T1 FS Cor Post T1 FS Sag Post			Perform T2 weighted image immediately after injection to allow for contrast uptake.	